



SHORT COMMUNICATION

Westernmost record of the diamondback puffer, *Lagocephalus guentheri* (Tetraodontiformes: Tetraodontidae) in the Mediterranean Sea: First record from Greek waters

Periklis Kleitou^{1,2,3} | Ioannis Giovos¹ | Francesco Tiralongo^{4,5} |
Nikolaos Doumpas¹ | Giacomo Bernardi⁶

¹iSea, Environmental Organization for the Preservation of the Aquatic Ecosystems, Greece

²Marine and Environmental Research (MER) Lab Ltd, Limassol, Cyprus

³School of Marine Science and Engineering, Plymouth University, Plymouth, UK

⁴Ente Fauna Marina Mediterranea, Avola, Italy

⁵Department of Biological, Geological and Environmental Sciences, University of Catania, Catania, Italy

⁶Department of Ecology and Evolutionary Biology, University of California Santa Cruz, Santa Cruz, California

Correspondence

Periklis Kleitou, iSea, Environmental Organization for the Preservation of the Aquatic Ecosystems, Greece.
Email: pkleitou@merresearch.com

Abstract

On 5th November 2017, a puffer fish specimen was caught in the Aegean Sea. Morphological and molecular analysis identified it as *Lagocephalus guentheri*. This first record from Greek waters represents the westernmost observation of the species in the Mediterranean Sea. The expansion and establishment of puffer fish species in the basin are discussed, with particular emphasis on the importance of the additional use of molecular analysis for increasing the efficiency in understanding species spread and dynamics.

1 | INTRODUCTION

The on-going climate change and human pressures in the Mediterranean Sea have led to substantial shifts of the species' communities, and expansion of tropical and subtropical species, particularly along the Eastern Mediterranean continental shelf which is susceptible to introductions via the Suez Canal (Arndt, Givan, Edelist, Sonin, & Belmaker, 2018; Givan, Edelist, Sonin, & Belmaker, 2018). In recent years, an expansion of pufferfish species (family Tetraodontidae) across the Mediterranean has been observed; raising concerns due to the accumulation of a potentially lethal neurotoxin in their tissues, and their potential impacts to the ecosystem, fisheries and economy (Giusti et al., 2018; Kalogirou, 2013). In this respect, carefully identifying the timing of their western advance in the Mediterranean is important, in order to better understand the dynamics of their invasion.

Lagocephalus is the most speciose genus, with five species recorded so far; namely, the native *L. lagocephalus* (Linnaeus, 1758), and four Indo-Pacific Lessepsian immigrants, *L. sceleratus* (Gmelin,

1789), *L. spadiceus* (Richardson, 1845), *L. suezensis* Clark & Gohar, 1953, and *L. guentheri* Miranda Ribeiro, 1915. *Lagocephalus* identification to species level is often ambiguous, leading to misidentifications and difficulties in allowing correct distribution assessments of the species in the Mediterranean. A strong example is the taxonomic issue that occurred with the misidentification of *Lagocephalus guentheri* Miranda Ribeiro, 1915 for its congeneric, *L. spadiceus* (Richardson, 1845). Their subtle distinguishing features led to several Mediterranean records of *L. spadiceus* to be questionable (Vella, Vella, Karakulak, & Oray, 2017). The misidentification issue has been addressed by Matsuura, Golani, and Bogorodsky (2011), who provided a clear description of the species, and identified morphological differences between the two species that are easy to score. Nevertheless, misidentifications may persist. Studies that utilize combined identification tools are likely to improve taxonomic knowledge and enable correct assessment of the species' spread in the basin (e.g. Farrag, El-Haweet, & Moustafa, 2016; Vella et al., 2017). This report presents the first record of *L. guentheri* in Greece, and the western-most record for the species in the Mediterranean basin.




FIGURE 1 (a) The red circle indicates the catch reported in this study, and the blue indicate confirmed sightings of *Lagocephalus guentheri* in the Mediterranean Sea; (b) Specimen analysed in this study

2 | MATERIALS AND METHODS

On 5th November 2017, a recreational fisherman caught a specimen of *L. guentheri* off the coasts of Pachi (Attiki) ($37^{\circ}57'22.7''\text{N}$ $23^{\circ}21'16.9''\text{E}$; Figure 1) at a depth of 24 m. The specimen was submitted in a citizen-science program and was collected for macroscopic analysis. Morphometric measurements were obtained and the specimen was identified based on macroscopic examination. In addition, DNA was extracted from the sample and the mitochondrial barcode gene CO1 (Cytochrome oxidase c subunit 1) was amplified following published protocols (Bariche et al., 2015). Briefly, the amplification of COI used fish specific primers VF2T1 and VR1dT1 (Ivanova, Zemlak, Hanner, & Hebert, 2007). PCR amplified fragments were sequenced in both directions using the same primers, and then compared with available sequences in GenBank. Phylogenetic reconstructions were performed based on the Neighbour-Joining method generated in R (R Core Team, 2016) with the use of the ape package (Paradis, Claude, & Strimmer, 2004). Genetic distances were calculated using a Kimura 2-parameter method. The maximum likelihood (ML) method has been also used as a second phylogenetic reconstruction approach, as implemented in GARLI (Zwickl, 2006). To estimate support for the nodes, 1,000 bootstrap replicates were

performed and we retained only the values supporting the nodes accounting for more than 50% of the bootstrap replicates.

3 | RESULTS

The specimen was identified as *L. guentheri* based on the morphological characteristics described by Matsuura et al. (2011). Specifically, pectoral fins were pale, while the caudal fin was slightly lunate, and an apparent posterior extension medially that made it appear to be doubly emarginated (Figure 1b); distinguishing it from its closest congener *L. spadiceus*. In addition, the caudal fin of the specimen was almost entirely dark-brown except of dorsal and ventral white tips, whereas in *L. spadiceus*, the ventral one-third of the caudal fin is white. The morphometric measurements collected are presented in Table 1.

The PCR amplification and sequencing of the cytochrome oxidase 1 resulted in a 652 bp fragment (GenBank accession number MH277032). A BLAST comparison of this sequence with available sequences in GenBank resulted in a 100% match with both *Lagocephalus spadiceus* and *L. guentheri*. In contrast, a query in BOLD resulted in a 100% positive identification to *L. guentheri*.

TABLE 1 Morphometric measurements of the *Lagocephalus guentheri* analysed in the present study

Total Length	23.9 cm
Fork Length	23.3 cm
Standard Length	20.6 cm
Head Length	6.4 cm
Pre-anal length	11.5 cm
Pre-pectoral length	5.7 cm
Body Depth	5.3 cm

Phylogenetic analysis was performed by comparing our sequence to 52 sequences extracted from GenBank (Figure 2). Misidentification of *Lagocephalus* individuals resulting in wrong taxonomic assignments of sequences is rampant. Vella et al. (2017) addressed this question and present convincing evidence as to the proper matches between good taxonomic individuals and their sequences. We therefore also included sequences that are considered as representative of *bona fide* species according to Vella et al. (2017), for *L. lagocephalus*, *L. spadiceus*, and *L. guentheri* from BOLD (boxed in red in Figure 2). Our sequence clustered with 30 GenBank sequence assigned to *L. spadiceus* (18 sequences), *L. guentheri* (10 sequences), and *Dactyloptena orientalis* (2 sequences) (Figure 2). The *bona fide*

L. guentheri species belonged to that cluster (although originally labeled *L. spadiceus*).

4 | DISCUSSION

Continuous, directed monitoring and management plan for the detection and abundance monitoring of alien species is imperative as the biodiversity of the Mediterranean basin is changing (Farrag et al., 2016). Inevitably, the addition of new tools such as genetics, enable the accurate taxonomic identification and taxonomic revision of the marine alien species in the Mediterranean as distributions, identifications and nomenclatures are updated. (Zenetos et al., 2017).

Vella et al. (2017) have pointed out that BOLD accession numbers AAD4510 and ADG5739 correspond to *L. spadiceus* and *L. guentheri*, respectively. The morphological analysis of our specimen is consistent with the characteristics of *L. guentheri*, and its COI sequence does cluster with the ADG5739 sequence showing concordance between morphological and genetic assignment of our specimen.

To date, the presence of *L. guentheri* (or individuals previously identified as *L. spadiceus*) remained confined to the Eastern Mediterranean and had not shown signs of westward expansion




FIGURE 2 Neighbour-Joining (NJ) phylogenetic tree based on K2P distances of mitochondrial cytochrome oxidase c subunit 1 sequences of *Lagocephalus* species (and identical general topology was obtained using Maximum Likelihood, ML). Bootstrap support values higher than 50% are shown next to the corresponding nodes and represent the NJ distance and ML value respectively. The sequence generated in this study (LGU Med) is highlighted in red. All other sequences are from GenBank. Three sequences that are considered as *bona fide* BOLD specimens are boxed in red. *L. spadiceus* and *L. guentheri* clades are boxed in green and yellow, respectively. Species native to the Mediterranean are boxed in blue and Indo-Pacific immigrants in red

(Azzurro, Maynou, Belmaker, Golani, & Crooks, 2016). Driven by the climate change and the synergistic effects of human pressures, it is possible that the species' dispersal, niche availability and establishment across the Mediterranean will be expanded on the following years. This study reports the westernmost record of the species in the Mediterranean basin. The improvement of taxonomic knowledge and utilization of identification tools that will incorporate morphological and molecular data should be facilitated to avoid common taxonomic issues of the past and monitor species' expansion to forestall any possible latent events.

ACKNOWLEDGMENTS

The authors are grateful to the fisherman Nikolaos Bacharakis for sharing the information and specimen to the citizen-science program, "Is it Alien to you.... Share it!!!", and to Eirini Gratsia for helping in the development of the map.

ORCID

Periklis Kleitou  <http://orcid.org/0000-0002-9168-4721>

Francesco Tiralongo  <http://orcid.org/0000-0002-1625-0149>

Giacomo Bernardi  <http://orcid.org/0000-0002-8249-4678>

REFERENCES

- Arndt, E., Givan, O., Edelist, D., Sonin, O., & Belmaker, J. (2018). Shifts in Eastern Mediterranean Fish Communities: Abundance changes, trait overlap, and possible competition between native and non-native species. *Fishes*, 3, 19. <https://doi.org/10.3390/fishes3020019>
- Azzurro, E., Maynou, F., Belmaker, J., Golani, D., & Crooks, J. (2016). Lag times in Lessepsian fish invasion. *Biological Invasions*, 18, 2761–2772. <https://doi.org/10.1007/s10530-016-1184-4>
- Bariche, M., Torres, M., Smith, C., Sayar, N., Azzurro, E., Baker, R., & Bernardi, G. (2015). Red Sea fishes in the Mediterranean Sea: A preliminary investigation of a biological invasion using DNA barcoding. *Journal of Biogeography*, 42, 2363–2373. <https://doi.org/10.1111/jbi.12595>
- Farrag, M., El-Haweet, A. A., & Moustafa, M. A. (2016). Occurrence of puffer fishes (Tetraodontidae) in the eastern Mediterranean, Egyptian coast-filling in the gap. *BiolInvasions Records*, 5, 47–54. <https://doi.org/10.3391/bir.2016.5.1.09>
- Giusti, A., Ricci, E., Guarducci, M., Gasperetti, L., Davidovich, N., Guidi, A., & Armani, A. (2018). Emerging risks in the European seafood chain: Molecular identification of toxic *Lagocephalus* spp. in fresh and processed products. *Food Control*, 91, 311–320. <https://doi.org/10.1016/j.foodcont.2018.04.013>
- Givan, O., Edelist, D., Sonin, O., & Belmaker, J. (2018). Thermal affinity as the dominant factor changing Mediterranean fish abundances. *Global Change Biology*, 24, e80–e89. <https://doi.org/10.1111/gcb.13835>
- Ivanova, N. V., Zemplak, T. S., Hanner, R. H., & Hebert, P. D. N. (2007). Universal primer cocktails for fish DNA barcoding. *Molecular Ecology Notes*, 7, 544–548. <https://doi.org/10.1111/j.1471-8286.2007.01748.x>
- Kalogirou, S. (2013). Ecological characteristics of the invasive pufferfish *Lagocephalus sceleratus* (Gmelin, 1789) in the eastern Mediterranean Sea—a case study from Rhodes. *Mediterranean Marine Science*, 14, 251–260. <https://doi.org/10.12681/mms.364>
- Matsuura, K., Golani, D., & Bogorodsky, S. V. (2011). The first record of *Lagocephalus guentheri* Miranda Ribeiro, 1915 from the Red Sea with notes on previous records of *L. lunaris* (Actinopterygii, Tetraodontiformes, Tetraodontidae). *Bulletin of the National Museum of Natural Science, Series A*, 37, 163–169.
- Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of phylogenetics and evolution in R language. *Bioinformatics*, 20, 289–290. <https://doi.org/10.1093/bioinformatics/btg412>
- R Core Team. (2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Retrieved from <https://www.R-project.org/>.
- Vella, A., Vella, N., Karakulak, F. S., & Oray, I. (2017). DNA Barcoding of Tetraodontidae Species from the Mediterranean Sea: Filling Knowledge Gaps for Improved Taxonomic Accuracy. *Genetics of Aquatic Organisms*, 1, 71–79. https://doi.org/10.4194/2459-1831-v1_2_05
- Zenetos, A., Çinar, M. E., Crocetta, F., Golani, D., Rosso, A., Servello, G., Verlaque, M. (2017). Uncertainties and validation of alien species catalogues: The Mediterranean as an example. *Estuarine, Coastal and Shelf Science*, 191, 171–187. <https://doi.org/10.1016/j.ecss.2017.03.031>
- Zwickl, D. (2006). GARLI: Genetic algorithm for rapid likelihood inference. Retrieved from <https://www.bio.utexas.edu/faculty/antisense/garli/Garli.html>.

How to cite this article: Kleitou P, Giovos I, Tiralongo F, Doumpas N, Bernardi G. Westernmost record of the diamondback puffer, *Lagocephalus guentheri* (Tetraodontiformes: Tetraodontidae) in the Mediterranean Sea: First record from Greek waters. *J Appl Ichthyol*. 2019;35:576–579. <https://doi.org/10.1111/jai.13811>