

Decapoda and Stomatopoda (Crustacea) of Rodos island (Greece) and the erythrean expansion NW of the Levantine sea

K. KEVREKIDIS¹ and B.S. GALIL²

¹ TEI Eperou, School of Fisheries & Aquaculture, 46 100 Igoumenitsa, Greece
e-mail: agrotiki @ionianislands.gr

² Israel Oceanographic & Limnological Research, National Institute of Oceanography,
P.O.B. 8030, Haifa 31080, Israel

Abstract

The Decapoda and Stomatopoda (Crustacea) of the sublittoral zone of Rodos island were studied between 1995 and 2000. Twenty-five sites, at depths ranging from the intertidal to 119 m, were sampled. Three species of stomatopods and 52 species of decapods were identified, bringing the number of decapods known from Rodos to 83. One Stomatopode and nineteen decapods, though previously known from the Aegean Sea, are new records for Rodos. The presence of seven species of Erythrean origin testify to a trend of increasing tropicalization of the area. A remarkable number of Erythrean taxa (fish, Crustacea, Mollusca, Polychaeta) have been established in Rodos, the Dodecanese and the southern Aegean Sea, whereas few reached the northern Aegean and the Ionian Seas. The expansion of the Erythrean species NW of the Levantine Sea is discussed.

Keywords: Crustacea, Decapoda, Stomatopoda, Rodos island, Aegean, Ionian, Levantine, Erythrean invasion.

Introduction

The study of the Crustacea of Rodos island has began in the early years of the 20th century (PARISI, 1913; COLOSI, 1923; SANTUCCI, 1928; MALDURA, 1938; TORTONESE, 1947a,b; KINZELBACH, 1965, 1970; PRETZMANN, 1971). Following the expedition of the Hebrew University - Smithsonian Institution to Rodos, LEWINSOHN (1976) published a list of the decapod crustaceans of Rodos. Eleven species of decapods were added

since (THESSALOU-LEGAKIS, 1986; THESSALOU-LEGAKI *et al.*, 1986; PANCUCCI-PAPADOPOULOU *et al.*, 1999; KOUKOURAS & DOUNAS, 2000). Only one stomatopod, *Squilla mantis* (L.), has been reported from Rodos (MALDURA, 1938; TORTONESE, 1947a,b).

A study of the sublittoral Decapoda and Stomatopoda of Rodos was conducted between 1995-2000. The material collected included 6 decapods and a stomatopod of Indo-Pacific origin (KEVREKIDIS & KEVREKIDIS, 1996; KEVREKIDIS *et al.*, 1998;

GALIL & KEVREKIDIS, 2002). A list of the entire decapod and stomatopod fauna of Rodos is presented.

Materials and Methods

Twenty-five sites along the coasts of Rodos island were sampled (Fig. 1), at depths ranging from the intertidal to 119 m. Samples were collected by a commercial trawler in deeper sites, and by diving in shallow ones. The samples are deposited in the School of Fisheries and Technology, Igoumenitsa, TEI Eperou. The site data (e.g. location, distance from shore, depth, sampling date) is presented in Table 1.

Results and Discussion

The island of Rodos, close to the coast of Asia Minor, borders the Aegean and the Levantine seas. The Rodos gyre, southeast of the island, and the Asia Minor Current (AMC)

are the major hydrological features nearby. The AMC runs along the Turkish coastline carrying westwards warm, salty water from the Levantine Sea, passing northward through the eastern Cretan Arc Straits, mainly the Rodos and Karpathos straits (THEOCHARIS *et al.*, 1993; NITTIS & LASCARATOS, 1999). Water masses of the area are characterised by high temperature and salinity values, even in deep waters (PANCUCCI-PAPADOPOULOU *et al.*, 1992). Surface temperature ranges between 17,6 °C in February and 26,4 °C in August and salinity is about 39‰ (NCMR, 1988). Since the hydrological characteristics of the area around Rodos resemble that of Levantine Sea, close faunal relations between the two were expected (SIMBOURA & NICOLAIDOU, 1993).

A total of 52 decapods were collected, increasing the number of decapod crustaceans known from Rodos to 83 (Table 2). Six species originated in the Indo-Pacific: *Marsupenaeus japonicus* (Bate, 1888), *Metapenaeopsis aegyptia* Galil 1990, *Metapenaeopsis mogiensis*




Fig. 1: Map of Rodos Island (Greece) with sampling stations.

Table 1
Code of sampling stations, sampling stations, day of sampling, distance from the coast and respective depth.

Code	Sampling stations	Date	Distance from coast (km)	Depth (m)
A1	Aerodromio	13.3.1995	0.74	29-31
A2	Genadi	23.3.1995	1.11-1.57	14,5-22
A3	Haraki	24.3.1995	1.48-1.85	14,5-40
A4	Kameiros	26.3.1995	1.7	47-64
B1	Pefkoi-Kiotari	3.10.1996	0.9-1.85	20-29
B2	Genadi-Pefkoi	3.10.1996	1.7-2	29-31
B3	Plimmiri-Genadi	3.10.1996	1.6-1.7	33-36.5
B4	Plimmiri-Genadi	4.10.1996	1.2-1.7	25,5-36,5
B5	Prassonisi-Plimmiri	4.10.1996	1.3-1.8	18-47,5
B6	Prassonisi-Karavolas	4.10.1996	1.4-1.5	29-33
C1	Plimmiri-M. Kavos	29.10.1997	1.5-3.3	33-51
C2	Prassonisi	29.10.1997	2.4-4.6	63-85
C3	Lahania-Plimmiri	30.10.1997	1.85	33
C4	Genadi-Kiotari	30.10.1997	1.67-2.2	31-49
C5	Pefkoi	30.10.1997	2.4-2.6	80,5-93,3
C6	Lardos-Kiotari	30.10.1997	0.93	18,3-20
D1	Isl. Strogillo-Isl. Ktenia	27.5.1998	0.93-3.05	49,4-69,5
E1	Isl. Ktenia-Mandilas	20.11.1999	1,3	29
F1	Isl. Strogillo-Isl. Ctenia	21.11.1999	4-4,6	109-119
F2	Isl. Strogillo-Isl. Ctenia	21.11.1999	0,9-2,4	29,7-40
F3	Glifada - Isl. Strogillo	21.11.1999	0,9-1,85	68-95
S1	Haraki	9.9.2000	-	0-1
S2	Lardos	9.9.2000	-	0-1
S3	Plimmiri	9.9.2000	-	0-1
S4	Archagellos	9.9.2000	-	0-1
S2	Lardos	10.9.2000	-	0-1

consobrina (Nobili, 1904), *Trachysalambria palaestinensis* (Steinitz, 1932), *Charybdis longicollis* Leene, 1938 and *Ixa monodi* Holthuis & Gottlieb, 1956 (KEVREKIDIS & KEVREKIDIS, 1996; KEVREKIDIS *et al.*, 1998; GALIL & KEVREKIDIS, 2002). Nineteen species, previously known from the Aegean sea (KOUKOURAS *et al.*, 1992; D'UDECEM D'ACQZ, 1999), are newly recorded from Rodos.

Three stomatopod species were identified from our samples: *Squilla mantis* (L., 1758), reported previously by MALDURA (1938) and TORTONESE (1947a,b); *Rissoides pallidus* (Giesbrecht, 1910), already reported from the Aegean sea (KOÇATAS & KATAGAN, 1995); and *Erugosquilla massavensis* (Kossmann, 1880), a species of Indo-Pacific origin (GALIL &

KEVREKIDIS, 2002). *E. massavensis* has been known along the Levantine coast, southern and southwestern Turkey, (Fethiye Bay, opposite the island of Rodos) (HOLTHUIS, 1961; KOÇATAS & KATAGAN, 1995; KOÇATAS, 1981) and from northern Kriti (Crete) (DOUNAS & STEUDEL, 1994).

TORTONESE (1947b) noted that 'The marine fauna [of Rodos] does not seem to be a very rich one', but that 'Crustaceans were fairly abundant'. A more recent study of the benthic macrofaunal communities off the NW coast of Rodos found that decapods were poorly represented (PANCUCCI-PAPADOPOULOU *et al.*, 1999). Those studies were far from exhaustive as 25 of the 52 decapod species collected during the present study are new records for Rodos. All but the six Indo-Pacific

Table 2
List of Stomatopoda and Decapoda (Crustacea) from Rodos island.

A. Stomatopoda												
	Maldura, 1938			Tortonese, 1947a,b			Present study		Sampling stations			
Squillidae												
<i>Erugosquilla massavensis</i> **							+		B1			
<i>Rissoides pallidus</i> *							+		C2			
<i>Squilla mantis</i>			+			+	+		C4			
B. Decapoda												
	Paris 1913	Colosi 1923	Santucci 1928	Maldura 1938	Tortonese 1947a,b	Kinzelbach 1965;1970	Pretzmann 1971	Lewinsohn 1976	Thessalou Legakis 1986; <i>et al</i> 1986;	Pancucci- Koukouras- Present Papadopoulou Dounas <i>et al.</i> 1999 2000	study	Sampling stations
Solenoceridae												
<i>Solenocera membranacea</i> *											+	C1,C2,C4, C5,D1
Penaeeidae												
<i>Marsupenaeus japonicus</i> **											+	A1,A2,A3 A4,B5,E1
<i>Melicertus kerathurus</i>					+	+						
<i>Metapenaepsis aegyptia</i> **											+	B1,B2,F2
<i>Metapenaepsis mogiensis consobrina</i> **											+	A3,B1,C6
<i>Parapenaeus longirostris</i> *											+	B2,C5
<i>Trachysalambria palaestinisensis</i> **											+	A3,B1
Sicyoniidae												
<i>Sicyonia carinata</i> *											+	C6
Pandalidae												
<i>Plesionika edwardsii</i>									+		+	C5,C6
<i>Plesionika narval</i>									+		+	B2
Alpheidae												
<i>Alpheus dentipes</i>									+			
<i>Syalpheus gambarelloides</i>						+						
Processidae												
<i>Processa acutirostris</i> *											+	B6,C2,C1
<i>Processa macrophthalma</i>											+	D1
Palaemonidae												
<i>Palaemon elegans</i>			+			+			+		+	S1,S2,S3,S4
<i>Palaemon xiphias</i>									+		+	B6
<i>Palaemonetes antennarius</i>									+			
<i>Pontonia pinnophylax</i>						+			+		+	C5

Table 2 (Continued)

B. Decapoda (cont.)														
	Parisi 1913	Colosi 1923	Santucci 1928	Maldura 1938	Tortonese 1947a,b	Kinzelbach 1965;1970	Pretzmann 1971	Lewinsohn 1976	Thessalou Legakis 1986; <i>et al.</i> 1986;	Pancucci- Papadopoulou <i>et al.</i> 1999	Koukouras- Dounas 2000	Present study	Sampling stations	
Crangonidae														
<i>Aegaon cataphractus*</i>												+	B1,C1,C2, C3,C4,C5,D1	
Palinuridae														
<i>Palinurus elephas</i>					+			+						
Scyllaridae														
<i>Scyllarides latus</i>				+	+								+	D1
<i>Scyllarus arctus</i>					+								+	B6
<i>Scyllarus pygmaeus</i>								+					+	C1
Callianassidae														
<i>Callianassa candida</i>								+					+	S2
<i>Callianassa subterranea</i>									+					
<i>Callianassa tyrrhena*</i>													+	S3
<i>Gouretia denticulata</i>									+		+			
Upogebiidae														
<i>Gebiacantha talismani</i>									+		+			
<i>Upogebia pusilla*</i>													+	S2,S3
<i>Upogebia stellata</i>									+		+			
<i>Upogebia tipica</i>								+	+		+			
Diogenidae														
<i>Calcinus tubularis</i>								+						
<i>Clibanarius erythropus</i>				+				+					+	S3
<i>Dardanus arrosor</i>					+								+	F3
<i>Dardanus calidus</i>								+					+	B6
<i>Diogenes pugilator</i>								+						
<i>Paguristes eremita</i>				+				+					+	C1
Paguridae														
<i>Anapagurus laevis</i>								+						
<i>Cestopagurus timidus</i>								+						
<i>Pagurus anachoretus</i>								+						
<i>Pagurus prideaux</i>								+					+	B3,C1
Galatheidae														
<i>Galathea intermedia</i>										+				
<i>Galathea machadoi</i>													+	
<i>Galathea squamifera</i>				+										
<i>Galathea strigosa</i>					+									
Porcellanidae														
<i>Pisidia bluteli</i>								+						
<i>Porcellana ptatycheles</i>								+						
Homolidae														
<i>Homola barbata*</i>													+	D1,F3

Table 2 (Continued)

B. Decapoda (cont.)												
	Parisi 1913	Colosi 1923	Santucci 1928	Maldura 1938	Tortonese 1947a,b	Kinzelbach 1965;1970	Pretzmann 1971	Lewinsohn 1976	Thessalou Legakis 1986; <i>et al</i> 1986;	Pancucci- Papadopoulou <i>et al.</i> 1999	Koukouras- Dounas 2000	Present Sampling study stations
Latreillidae												
<i>Latreillia elegans</i> *												+ F2
Dromiidae												
<i>Dromia personata</i>					+							+ D1
Dorippidae												
<i>Ethusa mascarone</i> *												+ F3
Calappidae												
<i>Calappa granulata</i>					+							+ B3,C3
Leucosiidae												
<i>Ilia nucleus</i> *												+ C6
<i>Ixa monodi</i> **												+ F2
Majidae												
<i>Acanthonyx lunulatus</i>			+		+			+				
<i>Eurynome aspera</i>										+		
<i>Inachus communissimus</i> *												+ B1,C4
<i>Inachus dorsettensis</i> *												+ D1,F2
<i>Inachus leptochirus</i> *												+ D1
<i>Inachus thoracicus</i> *												+ B2,C1, C4,F2
<i>Macropodia rostrata</i>										+		+ F2
<i>Maja crispata</i>	+				+			+				
<i>Maja squinado</i>					+							+ D1
<i>Pisa armata</i> *												+ C2,D1, F2,F3
<i>Pisa corallina</i>								+				+ B6
<i>Pisa muscosa</i> *												+ B6
<i>Pisa tetraodon</i>								+				
Portunidae												
<i>Callinectes sapidus</i>						+						
<i>Charybdis (Goniohellenus) longicollis</i> **												B1,C1,C3, + C4,C6,F2
<i>Liocarcinus arcuatus</i>			+		+							+ B6,F2
<i>Liocarcinus corrugatus</i>					+							+ B6,C2
<i>Liocarcinus depurator</i>					+							+ C2,D1
<i>Liocarcinus zariquieyi</i>								+				
<i>Portunus hastatus</i>					+							+ C1,C3, C4,C6
Xanthidae												
<i>Monodaeus guinotae</i> *												+ F1
<i>Xantho granulicarpus</i>								+				
<i>Xantho poressa</i>			+		+			+				+ S2,S3

Table 2 (Continued)

B. Decapoda (cont.)											
	Parisi	Colosi	Santucci	Maldura	Tortonese	Kinzelbach	Pretzmann	Lewinsohn	Thessalou	Pancucci-	Koukouras- Present
	1913	1923	1928	1938	1947a,b	1965;1970	1971	1976	Legakis 1986; <i>et al</i> 1986;	Papadopoulou <i>et al.</i> 1999	Dounas 2000 study stations
Eriphiidae											
<i>Eriphia verrucosa</i>		+	+		+						+ S1
Pilumnidae											
<i>Pilumnus hirtellus</i>				+							
Goneblacidae											
<i>Goneplax rhomboides</i>										+	
Grapsidae											
<i>Pachygrapsus marmoratus</i>			+		+		+	+			+ S2,S3
Ocypodidae											
<i>Ocypode cursor</i>							+				
Palicidae											
<i>Palicus caronii</i> *											+ C2

* New records for Rodos island previously known from the Aegean Sea

** Erythrean species

species (see above), occur in the Aegean Sea (KOUKOURAS *et al.*, 1992; D'UDEKEM D'ACQZ, 1999). Our results thus confirm BARASH & DANIN's (1989) statement that the fauna of Rodos shows biogeographical affinity with that of the Aegean Sea. The presence of the seven Erythrean species, however argues an Indo-Pacific incursion, and testify to a trend of increasing tropicalization of the area.

A considerable number of Erythrean invaders (fish, Crustacea, Mollusca, Polychaeta) have been established in the Levantine Sea, and some have spread into the southern Aegean Sea. Examination of the expansion of the Erythrean invaders NW of the Levantine Sea

shows that the majority of the species first appear at Rodos and the Dodecanese islands, and whereas a significant number spread into other parts of the southern Aegean Sea, few have reached the northern Aegean and the Ionian Seas (Table 3). This pattern of distribution is apparent for Erythrean fish as 'the prevailing currents direct the spreading of fishes along the Asiatic coasts northward and then westward toward the Aegean Islands' (PAPACONSTANTINO, 1990). Eight of the 16 Erythrean fish recorded from Rodos and the Dodecanese islands are known from the south Aegean (PAPACONSTANTINO, 1990; CORSINI & ECONOMIDIS, 1999; www.ciesm.org/atlas). No Erythrean fish has been reported

Table 3
Number of established Erythrean species of fish, Crustacea, Mollusca, and Polychaeta NW of the Levantine Sea.

	Fish	Crustacea (Decapoda-Stomatopoda)	Mollusca (Bivalvia-Gastropoda)	Polychaeta	Total
Rodos-Dodekanese islands	16	7	4	4	31
South Aegean Sea excl. Dodekanese	8	2	7	5	22
North Aegean Sea	-	-	-	5	5
Ionian Sea	5	-	2	2	9

from the north Aegean Sea, possibly due to unfavourable abiotic factors (PAPACONSTANTINO & TORTONESE, 1980), whereas 5 species have spread into the Ionian Sea (e.g., TORTONESE, 1967; KASPIRIS, 1976). Four Erythrean molluscs and four Erythrean polychaetes have been collected off Rodos (BARASH & DANIN, 1989; BEN-ELIAHU & FIEGE, 1996; www.ciesm/org/atlas), seven and five species respectively recorded in the south Aegean (FOUNTOULAKIS & SABELLI, 1999; SIMBOURA & NICOLAIDOU, 2001; www.ciesm/org/atlas), and two molluscs and two polychaetes have been found in the Ionian Sea (ARCIDIACONO & DI GERONIMO, 1976; SIMBOURA & NICOLAIDOU, 2001; www.ciesm/org/atlas). However, five polychaete species of Erythrean origin were found in the north Aegean sea (SIMBOURA *et al.*, 1995; ARVANITIDIS, 2000). The first Erythrean invader, *Siganus rivulatus* Forskål, 1775 was reported from the Dodecanese islands by BRUNELLI & BINI (1934) noting «è notevolmente diffuso e di facile cattura». Already TORTONESE (1947b) claimed that the fish fauna of Rodos acquired a tropical character, as some invading fish were so abundant they were exploited commercially (TORTONESE, 1947a). POR (1978) believed Rodos the farthest point Erythrean invaders would reach and rejected the possibility of them spreading into the Aegean Sea. However, spread they did: PAPACONSTANTINO (1990) suggested that inadequate sampling was at the root of the error. In the past decade a significant number of molluscs and polychaetes have been recorded in the south Aegean sea (VARDALA-THEODOROU, 1999; SIMBOURA & NICOLAIDOU, 2001).

Climatic factors play an important role in bioinvasion: POR (1990) maintained that the distribution of Erythrean immigrants 'will certainly expand or shrink according to the climatic evolution in the area' and that 'immigrants will expand further west if the climate warms up further' (POR, 1978), and

PAPACONSTANTINO (1990) attributed the westward advance of Erythrean invaders to the rise of the sea temperature in the '50s. Recently, GALIL & KEVREKIDIS (2002) attributed the influx of the exotic crustaceans into the southeastern Aegean Sea in the past decade to the augmented salinity and to the more extensive inflow of the AMC following the significant changes of the south Aegean water mass characteristics termed the Eastern Mediterranean Transient (EMT). The EMT provides favourable conditions for the maintenance and spread of Erythrean species in the south Aegean Sea and signals the possible invasion of the area by species of Levantine or Erythrean origin. Thus, if the present characteristics of the south Aegean hydrography persist, a modification of the composition and structure of the fauna is entirely possible.

Acknowledgements

The first author would like to thank Prof. A. Koukouras (AUTH) for his substantial contribution, Dr. M. Simbours (NCMR) for kindly providing information and Mr. D. Kazepis for sending me references (NCMR library). The same author would like also to thank Capt. J. Karidis and the crew of the trawler 'Panagia' in Rodos for their help all these years and Miss S. Tsitsirigou Tech. Ichthyology, for her help along the shores of the island.

References

- ARCIDIACONO, G. & DI GERONIMO, I., 1976. Studio biometrico di alcuni campioni di *Brahidontes variabilis* (Krauss). *Conchiglia*, 12: 61-74.
- ARVANITIDIS, C. 2000. Polychaete fauna of the Aegean Sea: new information. *Bull. Mar. Sci.*, 60(1): 73-96.
- BARASH, A. & DANIN, Z., 1989/90. Marine mollusca at Rhodes. *Israel J. Zool.*, 35: 1-74.
- BEN-ELIAHU, M. N. & FIEGE, D., 1996. Serpulid tube-worms (Annelida Polychaeta) of the Central

- and Eastern Mediterranean with attention to the Levant Basin. *Senckenb. Marit.*, 28 (1/3).
- BRUNELLI, G. & BINI, G., 1934. Sulla immigrazione di una specie di «Teuthis» dal Mar Rosso al Mar Egeo. *Atti Ac. Naz. Lincei*, 19:255-260.
- CIESM. Atlas of exotic species in the Mediterranean. (www.CIESM.org/atlas/)
- COLOSI, G., 1923. Crostacei Decapodi della Cirenaica. *Mem. R. Comitato Talassogr. Ital.* 104: 1-11.
- CORSINI, M. & ECONOMIDIS, P.S., 1999. Distribution extension of two Lessepsian migrants found in the marine area of the island of Rhodes (Aegean sea, Greece). *Cybium*, 23(2): 195-199.
- DOUNAS, C. & STEUDEL, C., 1994. Stomatopod Crustacea from the island of Crete. *Crustaceana*, 66(2): 252-254.
- FOUNTOULAKIS, E. & SABELLI, B., 1999. Observations on the malacofauna of the littoral of Saronikos Gulf. In: *Contributions to the zoogeography and ecology of the Eastern Mediterranean region*, vol. 1: 465-472.
- GALIL, B.S. & KEVREKIDIS, K., 2002. Exotic decapods and stomatopod off Rhodes (Rodos)(Greece) and the Eastern Mediterranean Transient. *Crustaceana*, 75(7): 925-930.
- HOLTHUIS, L.B., 1961. Report on a collection of Crustacea Decapoda and Stomatopoda from Turkey and the Balkans. *Zool. Verhand.*, 47: 1-67.
- KASPIRIS, P., 1976. New fish records from the Greek part of the Ionian Sea. *Rev. Trav. Inst., Peches marit.*, 40 (3 et 4): 627-628.
- KEVREKIDIS, K. & KEVREKIDIS, T., 1996. The occurrence of *Penaeus japonicus* Bate (Decapoda, Penaeidae) in the Aegean sea. *Crustaceana*, 69 (7): 925-929.
- KEVREKIDIS, K., GALIL, B. S. & KEVREKIDIS, T., 1998. Three Lessepsian migrant penaeids (Decapoda) in Rodos island (Greece). *Crustaceana*, 71 (4): 474-478.
- KINZELBACH, R., 1965. Die Blaue Schwimmkrabbe (*Callinectes sapidus*) ein Neuburger im Mittelmeer. *Nat.Mus. (Frankfurt)* 95: 293-296.
- KINZELBACH, R., 1970. Neue Nachweise der Reiterkrabbe, *Ocypode cursor* (Linnaeus, 1758), in der Agais (Decapoda, Brachyura, Ocypodidae). *Crustaceana*, 18(3): 318-320.
- KOÇATAS, A., 1981. Two new stomatopod species for the Turkish fauna: *Nannosquilla loides* occultus (Giesbrecht) and *Meiosquilla desmaresti* (Riso). *Crustaceana*, 40 (2): 318-320.
- KOÇATAS, A. & KATAGAN, T., 1995. On Stomatopoda from Turkey with the first record of *Rissoides pallidus* for the Turkish fauna. *Crustaceana*, 68(5): 649-652.
- KOUKOURAS, A. & DOUNAS, C., 2000. Decapod Crustaceans new to the fauna of the Aegean sea. *Crustaceana*, 73(4): 497-502.
- KOUKOURAS, A., DOUNAS, C., TURKAY, M. & VOULTSIADOU-KOUKOURA, E., 1992. Decapod crustacean fauna of the Aegean sea: new information, check list, affinities. *Senckenb. marit.*, 22: 217-244.
- LEWINSOHN, CH., 1976. Crustacea Decapoda von der Insel Rhodos, Griechenland. *Zool. Meded. (Leiden)*, 49: 237-254.
- MALDURA, C. M. 1938. La pesca nelle Isole Italiane dell' Egeo. *Boll.pesca piscic. Idrobiol.* 14: 460-481.
- NATIONAL CENTRE FOR MARINE RESEARCH (N.C.M.R.), 1988. Oceanographic and ecological survey of the SE Aegean Sea (1983-84). Technical Report, Athens (in Greek).
- NITTIS, K. & LASCARATOS, A., 1999. Intermediate water formation in the Levantine sea: the response to interannual variability of atmospheric forcing. In: *The Eastern Mediterranean as a laboratory basin for the assessment of contrasting ecosystems*, edited by P. MALANOTTE-RIZZOLI and V. N. EREMEEV, Kluwer Academic Publishers, Dordrecht: 441-446.
- PANCUCCI-PAPADOPOULOU, M. A., SIOKOU-FRANGOU, I., THEOCHARIS, A. & GEORGOPOULOS, D., 1992. Zooplankton vertical distribution in relation to the hydrology in the NW Levantine and the SE Aegean seas (spring 1986). *Oceanologica Acta*, 15: 365-381.
- PANCUCCI-PAPADOPOULOU, M. A., SIMBOURA, N., ZENETOS, A., THESSALOU-LEGAKI, M. & A. NICOLAIDOU, 1999. Benthic invertebrate communities of NW Rodos (Rhodes) island (SE Aegean Sea) as related to hydrological regime and geographical location. *Isr. J. Zool.*, 45: 371-393.
- PAPACONSTANTINO, C., 1990. The spreading of lessepsian fish migrants into the Aegean Sea (Greece). *Sci. Mar.*, 54(4): 313-316.
- PAPACONSTANTINO, C. & TORTONESE, E., 1980. On a collection of fishes from the Thermaikos Gulf (N.E. Greece). *Thalassografica*, 3(2): 15-42.

- PARISI, B., 1913. Decapodi. Escursioni Zoologiche del Dr. Enrico Festa nell' Isola di Rodi. *Boll. Musei Zool. Anat. comp. R. Univ. Torino*, 28: 1,2.
- POR, F. D., 1978. Lessepsian migration. The influx of Red Sea biota into the Mediterranean by way of the Suez Canal. *Ecological studies*, 23, Springer, Berlin.
- POR, F. D., 1990. Lessepsian migration. An appraisal and new data. *Bulletin de l'Institut oceanographique*, Monaco, n° special 7: 1-10.
- PRETZMANN, G., 1971. Ergebnisse einiger Sammelreisen nach Vorderasien, 2. Teil: Marine Brachyura. *Ann. Naturhistor. Mus. Wien*, 75: 477-487.
- SANTUCCI, R., 1928. Alcuni Crostacei Decapodi delle isole Egee. Ricerche faunistiche nelle isole italiane dell' Egeo. *Arch. zool. Ital.*, 12:345-354.
- SIMBOURA, N. & NICOLAIDOU, A., 1993. The polychaete fauna of the northern coast of Rhodes and comparison with other areas. *Proc. 4th Natl. Symp. Oceanogr. Fish.*, Rodos: 210-213 (in Greek).
- SIMBOURA, N. & NICOLAIDOU, A., 2001. The polychaetes (Annelida, Polychaeta) of Greece: checklist, distribution and ecological characteristics. *Monographs on Marine Sciences, Series no 4*, NCMR, Athens.
- SIMBOURA, N., ZENETOS, A., THESSALOU-LEGAKI, M., PANCUCCI, M. A. & NICOLAIDOU, A., 1995. Benthic communities of the infralittoral in the N. Sporades (Aegean sea): a variety for biotopes encountered and analysed. *P.S.Z.N.I., Marine Ecology*, 16(4): 283-306.
- THEOCHARIS, A., GEORGOPOULOS, D., LASCARATOS, A. & NITTIS, K., 1993. Water masses and circulation in the central region of the Eastern Mediterranean: Eastern Ionian, South Aegean and Northwest Levantine, 1986-1987. *Deep-Sea Res. II. Top. Stud. Oceanogr., Spec. Issue*, 40: 1121-1142.
- THESSALOU-LEGAKIS, M., 1986. Preliminary data on the occurrence of Thalassinidae (Crustacea: Decapoda) in the Greek seas. *Biol. Gallo-Hell.*, 12: 181-187.
- THESSALOU - LEGAKI, M., FRANTZIS, A., NASSIOKAS, K. & HATZINIKOLAOU, S., 1986. Observations on *Parapandalus narval* (Fabricius, 1787) (Crustacea, Decapoda, Pandalidae) from Rodos island (Greece). *Rapp. Comm. Int. Mer medit.*, 30, 2.
- TORTONESE, E., 1947a. Note intorno alla fauna e flora marine dell' Isola di Rodi (Mar Egeo). *Boll. Pesca Piscic. Idrobiol.*, 23: 13-20.
- TORTONESE, E., 1947b. Biological investigations in the Aegean sea. *Nature*, London 159: 887.
- TORTONESE, E., 1967. Un pesce plettognato nuovo per I mari Italiani: *Stephanolepis diaspros*, Fraser Brunner. *Doriana, suppl. Annali der Museo civico di storia naturale di Genova*, 4 (181) : 1-4.
- UDEKEM D'ACCOZ, C. D'., 1999. Inventaire et distribution des Crustaces Decapodes de l' Atlantique nord-oriental, de la Mediterranee et des eaux continentales adjacentes au nord de 25°. N. Patrimoines Naturels (M.N.H.N./S.P.N.), 40:1-383 (Mus. Nat. d'Histoire Naturelle, Paris).
- VARDALA-THEODOROU, G.E., 1999. The occurrence of the Indo-Pacific molluscan species *Fulvia fragilis* (Forsskal, 1775) and *Bulla ampulla* L. 1758 in Elefsis Bay. Newsletter of the Hellenic Zoological Society, Fasc. 31 : 10-11

Dedication

Sailing near to Asia Minor; motherland of both my grandmother Alexandra KEVREKIDOU (Kars, Pontos) and my grandfather Dimitrios VOGDANOU (Proussa, Aeolia). A need for dedication of this work to their memory.