

BIOLOGICAL SURVEY OF THE EASTERN MEDITERRANEAN SEA : HYDROIDS
(PRELIMINARY STUDY).

J. MARINOPOULOS

Station marine d'Endoume 13007 - Marseille, France.

Résumé : Une importante collection d'Hydroïdes a été récoltée lors de l'expédition du N/O "Calypso" (1977) ; des 50 prélèvements réalisés jusqu'à 50 mètres de profondeur, 45 espèces d'Hydroïdes ont été déterminées ; toutes ont une large répartition méditerranéenne à l'exception de 2 espèces, restreintes au seul bassin oriental.

The biological survey of the eastern Mediterranean Sea was started in autumn 1977 during the "Calypso" cruise.

The survey involved qualitative and quantitative samplings of the benthos on hard-substrates to identify the species occurring in this region and to provide informations concerning their ecological requirements.

The hydroid fauna of the eastern Mediterranean Sea is poorly documented in comparison with that of other regions. Few studies were published on eastern Mediterranean Sea hydroids, based on small collections from a restricted area and without any ecological information.

A rich and diversified hydroid fauna occurs in the eastern Mediterranean sea : using scuba-diving and underwater photography up to 50 meters deep ; 50 samples were performed, each provided with numerous ecological observations. Forty-five species of hydroids were identified ; fourteen occurred in anfractuosités at the upper level of the "étage infralittoral" or sampled in deeper bottoms. Only thirteen are in strictly photophilic position.

The hydroids have a wide distribution in the Mediterranean Sea excepted for two species, *Aglaophenia* cf. *elongata* and *Sertularia marginata* which are restricted to the eastern basin.

Stations	TURQUIE								GRECE	
	1		2		3		4		5	
	P	S	P	S	P	S	P	S	P	S
<i>Halocordyle disticha</i>	+	.	.	.
<i>Eudendrium ramosum</i>	+	+	+	+	.	+	+	.	+	.
<i>E. glomeratum</i>	+
<i>E. motzkossowskiae</i>	+	+
<i>E. racemosum</i>	+	+	+
<i>E. capillare</i>	+	+	.
<i>Cordylophora neapolitana</i>	+	.
<i>Podocoryna fucicola</i>	.	+
<i>Garveia grisea</i>	.	+
<i>Laodicea undulata</i>	.	.	.	+	.	+
<i>Hobella scandens</i>	.	+
<i>H. parasitica</i>	.	+	.	+
<i>Scandis pocillum</i>	+
<i>Orthopyxis alta</i>	+
<i>O. crenata</i>	.	.	+
<i>O. caliculata</i>	+	.	+
<i>Clytia uniflora</i>	+	+	+	+	.	+	+	.	.	+
<i>C. gravieri</i>	+	+	.	+	+	+
<i>C. paulensis</i>	.	.	.	+
<i>Obelia geniculata</i>	+
<i>Laomedea calceolifera</i>	+	+
<i>Halecium mediterraneum</i>	.	.	.	+
<i>H. banyulense</i>
<i>E. pusillum</i>
<i>H. nanum</i>	+
<i>H. lankesteri</i>	+	.	+
<i>Kirchenpaueria pinnata</i>	+
<i>K. echinulata</i>	+
<i>K. echinulata f. similis</i>	+	+
<i>K. echinulata f. minuta</i>
<i>Plumularia setacea</i>	.	+	.	+
<i>Monothecha obliqua</i>	.	.	+	+
<i>M. posidoniae</i>
<i>Polyplumaria diaphana</i>	+	.	.	+
<i>P. secundaria</i>	.	+	+
<i>P. catharina</i>	+
<i>Aglaophenia octodonta</i>	+	+	+	+
<i>A. cf. elongata</i>	+	+	+	+
<i>Sertularella ellisii</i>	.	+	+	+	.	+
<i>S. polyzonias</i>	.	+	+
<i>Dynamena disticha</i>	+	+	+	+	.	.	+	.	+	.
<i>D. desmoides</i>	.	+	.	+
<i>Sertularia perpusilla</i>	+
<i>S. marginata</i>	.	.	+
<i>Fillelum serratum</i>	+

(P = photophilous ; S = sciaphilous).

(1 = Dorian Promontory ; 2 = Injeh's Cap ; 3 = Ismir Bay ;
4 = Antalya ; 5 = Aspra Spitia).