

First record of the New Zealand mud snail *Potamopyrgus antipodarum* J.E. Gray 1843 (Mollusca: Hydrobiidae) in Greece – Notes on its population structure and associated microalgae

Canella Radea, Ioanna Louvrou and Athena Economou-Amilli*

Department of Ecology & Systematics, Faculty of Biology, University of Athens, Panepistimiopolis, GR 157 84, Athens, Greece

*Corresponding author

E-mail: aamilli@biol.uoa.gr

Received 7 February 2008; accepted in revised form 30 July 2008; published online 5 October 2008

Abstract

Potamopyrgus antipodarum (Gastropoda: Hydrobiidae), a New Zealand native species, is recorded for the first time in waterbodies of Greece, being dominant in the gastropod fauna present; waterfowl transportation is suggested as its invasion mode. This species was collected (November 2007) from the aquatic plants *Myriophyllum spicatum* and *Nasturtium officinale* in Lake Trichonis (CW Greece) and an adjacent stream. Totally, 332 individuals of various ages were recorded, females being the majority. One diatom (*Cocconeis placentula* var. *euglypta*) was found in its digestive tube, also thriving abundantly on the surface of both the invader shell and the aquatic plants, and in association with other cosmopolitan periphytic diatoms (three of them new for the area, with one new record for Greece) or endemic diatoms.

Key words: alien species, Hydrobiidae, *Potamopyrgus antipodarum*, epizoic and periphytic diatoms, Lake Trichonis

Potamopyrgus antipodarum (J.E. Gray 1843) (Gastropoda: Hydrobiidae) is a snail species indigenous to New Zealand, which was introduced into Europe at the end of 19th century (Ponder 1988). In Europe, it is the most widespread non-indigenous prosobranch species and only a few countries have escaped invasion (Bank 2007), although its presumed absence could be due to the lack of recent field observations (Cianfanelli et al. 2007). In the Mediterranean basin, this species was recorded in Catalonia in 1952, in the Mediterranean region of France by the end of fifties (Falniowski 1987), and in Italy in 1961 (Cianfanelli et al. 2007). Shells of this species have also been found in Turkey (Demir 2003; Yildirim et al. 2006). No


previous data exist on the presence of this invader in waterbodies of Greece (Bank 2007).

P. antipodarum shows a considerable variability in shell dimensions (Falniowski 1987), its shell height reaching ~ 6.5 mm (Falniowski 1987; Richards et al. 2002). Being able to reproduce parthenogenically, females prevail (Falniowski 1987; Zaranko et al. 1997) and populations densities are occasionally extremely high (Kerans et al. 2005). *P. antipodarum* is classified as a scraper/grazer (Richards et al. 2002). It feeds on plant and animal detritus and it also grazes on green algae and diatoms, with valves of the latter group being found into its intestinal tube (Michaut 1968; Haynes and Taylor 1984; Kelly and Hawes 2005). It has also

been reported that the surface of its shell is often encrusted and covered with sediment and periphytic micro-organisms (Falniowski 1987; Richards et al. 2002).

In November 2007 during a student field excursion, *P. antipodarum* specimens were found in central-western Greece: a) in the littoral zone of Lake Trichonis (38°36'00"N, 21°34'11"E), a warm monomictic, rather oligotrophic lake (Overbeck et al. 1982; Tafas and Economou-Amilli 1997), and b) in a permanent, slowly flowing stream discharging into Trichonis near Dougri (38°36'01"N, 21°34'13"E). Samples from

all dominant aquatic vascular plants were taken (500g of plant material per species) using a grab sampler in Lake Trichonis at 1-3 m depth (with the kind help of Dr. D. Danielidis) or removed by hand from the stream. The snails occurred on macrophytes (kindly identified by Dr. I. Bazos), specifically only on the shoots of *Myriophyllum spicatum* L. (lake) and on the roots of *Nasturtium officinale* R. Br. (stream). Totally, 332 individuals of *P. antipodarum* of both genders and of various ages were collected (Figures 1-5) from the above plant species, making up 57% of the gastropod fauna present.




Figures 1-3. Specimens of *P. antipodarum* (1: immatures; 2-3: matures). Stereomicrographs. A background square represents 1 mm². Figure 4. Head with penis (eye at arrow). Stereomicrograph. Scale bar=1 mm. Figure 5. Penis (eye at arrow). Scanning Electron Micrograph (SEM). Scale bar=100 μ m. Figure 6. General view of attached diatoms on the shell surface of *P. antipodarum*. SEM. Scale bar=100 μ m. (Photographs by I. Louvrou and C. Radea).

The shell length of the specimens was up to 3mm. Females constituted the majority of the mature individuals, comprising approximately 78% of dissected specimens with 4 and/or 5 whorls.

The intestinal tubes' contents of 20 dissected individuals of *P. antipodarum* were also examined and valves of the diatom *Cocconeis placentula* var. *euglypta* were found. This diatom variety (Figure 14) abundantly colonizes, as epizoic taxon on the shell surface of these gastropods (Figure 6) and as periphytic taxon on the external surface of the aquatic plants (*M. spicatum* and *N. officinale*). Among the accompanying algal species present we recorded the scattered presence of the known periphytic (Krammer and Lange-Bertalot 1986, 1991a, 1991b) diatoms *Ulnaria ulna* (Nitz.) Compère, *Planothidium lanceolatum* (Bréb.) Round et Bukht., *P. frequentissimum* (Lange-Bert.) Round et Bukht., *Amphora pediculus* (Kütz.) Grunow, *Gomphonema angustatum* (Kütz.) Rabh. (Figures 7, 10-13); also *Cyclotella trichonidea* Ec.-Am. (Figure 8) and *C. trichonidea* var. *parva* Ec.-Am. (Figure 9), endemic (Economou-Amilli 1979, 1982) planktonic and periphytic (Economou-Amilli and Tafas 2000) diatoms. It is worth mentioning, that in Lake Trichonis, *Ulnaria ulna* and *C. placentula* var. *euglypta* have already been found as epizoic diatoms on another gastropod, i.e. on *Valvata piscinalis* O.F. Müll. (Falniowski et al. 1988); in addition, *Planothidium lanceolatum* was also found there as a planktonic diatom (Tafas and Economou-Amilli 1997). The other three diatom taxa are considered as new records for Lake Trichonis, and in fact, *Planothidium frequentissimum* as a new record for Greece.

P. antipodarum is distributed in most European countries (Bank 2007) and in Turkey (Aegean Sea, and Sea of Marmara) (Demir 2003; Yildirim 2006), but it has not been reported from the Balkan peninsula (Bank 2007) except for the Romanian Danube (Cioboiu 2006). Although the invasion mode of *P. antipodarum* in the wider area of Lake Trichonis still remains unknown, we suggest that it has possibly been transported from Europe and Turkey by resting and/or wintering waterfowl. It should be mentioned that Lake Trichonis is included in the Important Bird Areas of Greece (BirdLife International 2008) since numerous migratory waterfowl - many of them protected species - are recorded there every year. According to the existing information,



Figures 7-14. Epizoic and epiphytic diatoms after a light (by burning on a coverslip) valve cleaning treatment. 7. *Ulnaria ulna*, 8. *Cyclotella trichonidea*, 9. *C. trichonidea* var. *parva*, 10. *Gomphonema angustatum*, 11. *Planothidium lanceolatum*, 12. *P. frequentissimum*, 13. *Amphora pediculus*, 14. *Cocconeis placentula* var. *euglypta*. Light Micrographs (LM). Scale bar=10µm. (Photographs by I. Louvrou and A. Economou-Amilli).

P. antipodarum is able to withstand desiccation several days and is transported long distances by birds adhered to their bills and legs (Falniowski 1987; Global Invasive Species Database 2008; Bersine et al. 2008).

High abundance of *P. antipodarum* in studied Greek water bodies indicates an instant dominance in the indigenous gastropod community. In some cases, presence of the invader *P. antipodarum* may have a negative influence on native macroinvertebrates (Kerans et al. 2005), especially in disturbed ecosystems (Strzelec 2005). In other cases, there is evidence of a positive influence between densities of *P. antipodarum* and of some native benthic invertebrate species (Schreiber et al. 2002). Further investigations in the wider area of Lake Trichonis

should clarify the influence of this invader on the indigenous benthic community structure and on the ecological processes related to it.

References

- Bank RA (2007) Mollusca: Gastropoda. Fauna Europea version 1.3 (19 April 2007). <http://www.faunaeur.org/>. Cited 25 Feb 2008.
- Bersine K, Brenneis V, Draheim R, Rub AW, Zamon JE, Litton K, Hinton SA, Sytsma MD, Cordell JR and Chapman JW (2008) Distribution of the invasive New Zealand mudsnail (*Potamopyrgus antipodarum*) in the Columbia River Estuary and its first recorded occurrence in the diet of juvenile Chinook salmon (*Oncorhynchus tshawytscha*). Biological Invasions DOI10.1007/s10530-007-9213-y
- BirdLife International (2008) European Important Bird Areas. <http://www.birdlife.org>. Cited 26 Feb 2008
- Cianfanelli S, Lori E and Bodon M (2007) Non-indigenous freshwater molluscs and their distribution in Italy. In: Gherardi F (ed) Biological invaders in inland waters: Profiles, distribution, and threats, Springer, Dordrecht, The Netherlands, pp 103-121
- Ciobiou O (2006) Diversity of Gastropoda in the Romanian sector of the Danube lower hydrographic basin In: Proceedings 36th International Conference of IAD. Austrian Committee Danube Research / IAD, Vienna, 4-8 September 2006
- Demir M (2003) Shells of Mollusca Collected from the Seas of Turkey. Turkish Journal of Zoology 27: 101-140
- Economou-Amilli A (1979) Two New Taxa of Cyclotella from Lake Trichonis, Greece. Nova Hedwigia, 31: 467-477
- Economou-Amilli A (1982) SEM-Studies on *Cyclotella trichonidea* (Bacillariophyceae). Archiv für Hydrobiologie, Suppl. 63 (1), Algological Studies 30: 25-34
- Economou-Amilli A and Tafas T (2000) Distribution patterns of *Cyclotella trichonidea* Ec.-Am. sensu lato in the plankton succession of the lakes Trichonis and Amvrakia, Greece. Archiv für Hydrobiologie 147: 559-575
- Falniowski A (1987) Hydrobioidea of Poland (Prosobranchia: Gastropoda). Folia Malacologica 1: 1-122
- Falniowski A, Economou-Amilli A and Anagnostidis K (1988) *Valvata piscinalis* O.F. Müller (Mollusca-Prosobranchia) and its epizoic diatoms in Lake Trichonis, Greece. Internationale Revue der gesamten Hydrobiologie 73: 327-335
- Global Invasive Species Database (2008) *Potamopyrgus antipodarum* <http://www.issg.org/database/species/ecology.asp?si=449&fr=1&sts=> Cited 26 Feb 2008
- Haynes A and Taylor BJR (1984) Food finding and food preference in *Potamopyrgus jenkinsi* (E.A. Smith) (Gastropoda: Prosobranchia). Archiv für Hydrobiologie 100 (4): 479-491
- Kelly DJ and Hawes I (2005) Effects of invasive macrophytes on littoral zone productivity and foodweb dynamics in a New Zealand high-country lake. Journal of the North American Benthological Society 24: 300-320
- Kerans BL, Dybdahl MF, Gangloff MM and Jannot JE (2005) *Potamopyrgus antipodarum*: distribution, density, and effects on native macroinvertebrate assemblages in the Greater Yellowstone Ecosystem. Journal of the North American Benthological Society: Vol. 24 (1): 123-138
- Krammer K and Lange-Bertalot H (1986) Teil: Naviculaceae. In: Ettl H, Gerloff J, Heyning H and Mollenhauer D (eds) Süßwasserflora von Mitteleuropa. Bacillariophyceae 1. Fischer, Stuttgart, 876pp
- Krammer K and Lange-Bertalot H (1991a) Teil: Centrales, Fragilariaceae, Eunotiaceae. In: Ettl H, Gerloff J, Heyning H & Mollenhauer D (eds) Süßwasserflora von Mitteleuropa Bacillariophyceae 3. Fischer, Stuttgart, 576pp
- Krammer K and Lange-Bertalot H (1991b) Teil: Achnantheaceae, Kritische Ergänzungen zu *Navicula* (Lineolatae) und *Gomphonema*. In: Ettl H, Gerloff J, Heyning H & Mollenhauer D (eds) Süßwasserflora von Mitteleuropa. Bacillariophyceae 4. Fischer, Stuttgart, 437pp
- Michaut P (1968) Données biologiques sur un Gasteropode Prosobranch récemment introduit en Côte-d' Or, *Potamopyrgus jenkinsi*. Hydrobiologia 32:513-527
- Overbeck J, Anagnostidis K and Economou-Amilli A (1982) A limnological survey of three Greek lakes: Trichonis, Lyssimachia and Amvrakia. Archiv für Hydrobiologie 95 (1): 365-394
- Ponder WF (1988) *Potamopyrgus antipodarum*: a molluscan colonizer of Europe and Australia. Journal of Molluscan Studies 54: 271-285
- Richards D, Kerans B and Gustafson D (2002) New Zealand Mudsnail in the Western USA. Montana State University: Department of Ecology. <http://www.esg.montana.edu/aim/mollusca/nzms>. Cited 26 Feb 2008
- Schreider ESG, Lake PS and Quinn GP (2002) Facilitation of native stream fauna by an invading species? Experimental investigations of the interaction of the snail *Potamopyrgus antipodarum* (Hydrobiidae) with native benthic fauna. Biological Invasions 4: 317-325
- Strzelec M (2005) Impact of the introduced *Potamopyrgus antipodarum* (Gastropods) on the snail fauna in post-industrial ponds in Poland. Biologia (Bratislava) 60(2):159-163
- Tafas TP and Economou-Amilli A (1997) Limnological survey of the warm monomictic lake Trichonis (central western Greece). II. Seasonal phytoplankton periodicity – a community approach. Hydrobiologia 344: 141-153
- Yildirim MZ, Koca SB and Kebapçı U (2006) Supplement to the Prosobranchia (Mollusca: Gastropoda) Fauna of Fresh and Brackish Waters of Turkey. Turkish Journal of Zoology 30: 197-204
- Zaranko DT, Farara DG and Thompson FG (1997) Another exotic mollusc in the Laurentian Great Lakes: the New Zealand native *Potamopyrgus antipodarum* (Gray 1843) (Gastropoda, Hydrobiidae) Canadian Journal of Fisheries and Aquatic Sciences 54: 809-814